


Anti-Listeria Activity of Bacteriocin Produced by *Pediococcus pentosaceus* ST0408, a Potential Bio-Protective Culture

Kayque Ordonho Carneiro¹, Joao Marcos Scafuro Lima¹, Katia Leani¹, SVETOSLAV DIMITROV TODOROV^{1,2}

¹. Faculdade de Ciências Farmacêuticas, Universidade de São Paulo, ProBacLab, Laboratório de Microbiologia de Alimentos, Departamento de Alimentos e Nutrição Experimental, Sao Paulo, Brazil

². Instituto Politécnico de Viana do Castelo, CISAS - Center for Research and Development in Agrifood Systems and Sustainability, Viana Do Castelo, Portugal

One of the well-studied antimicrobial compounds produced by lactic acid bacteria (LAB) are bacteriocins, which are defined as polypeptides, produced via ribosomes and are typically antagonistic versus organisms that are genetically similar to the producers. Strains belonging to the species *Pediococcus pentosaceus* were associated to various fermented food products and have role in production of bacteriocins, metabolites involve in the food biopreservation. However, safety assessment and screening for presence of virulence and antibiotic resistance genes in LAB is an important task in the process for safety evaluation and suggestion for applications as commercial beneficial cultures. The aim of this study was to explore bacteriocinogenic and safety features of *P. pentosaceus* ST0408 based on biochemical, physiological antagonistic properties and presence and expression of genes related to the virulence factors, production of biogenic amines and antibiotic resistance. Bacteriocinogenic *P. pentosaceus* ST0408 was isolated from Bulgarian white brine cheese and identified based on its biochemical and genetic characteristics including 16S rRNA sequencing. Bacteriocin production was evaluated, including stability of antimicrobial peptide in different levels of pH (2.0-10.0), temperatures (10°C-121°C), presence of various chemicals applied in food processing processes. Activity of the expressed bacteriocin activity was tested versus numerous enterococci, lactobacilli, and listeria strains. Bacterial growth was monitored based on the changes culture turbidity (OD at 600 nm spectrophotometrically when *P. pentosaceus* ST0408 was cultured in MRS broth at 37°C for 24h; changes in pH were monitored; and production of bacteriocin was evaluated versus different strains of *L. monocytogenes*. Moreover, produced by *P. pentosaceus* ST0408 bacteriocins was added to exponentially growing *L. monocytogenes* strains and evaluated killing properties of expressed antimicrobials over 12h period. Produced by *P. pentosaceus* ST0408 bacteriocin was showing stable when exposed to the investigated pH levels, temperatures and chemicals used in dairy industry. Based on the results from performed biochemical and physiological test, such as hemolytic, gelatinase, proteolytic and lipolytic activity, biogenic amines and antibiotic resistance, *P. pentosaceus* ST0408 can be considered as safe strain. Moreover, strain ST0408 did not show evidence for presence of virulence genes, including vanA, B, C, D, E and G. *P. pentosaceus* ST0408 was expressing 51200 AU/mL bacteriocin activity recorded versus *L. monocytogenes* L211 during stationary growth phase. The addition of 6400 AU/mL of studied bacteriocin (pH 6.0) to a 3-h-old culture of *L. monocytogenes* L211 (OD 600nm 0.274) resulted in growth inhibition for monitored 12 h, suggesting that the mode of activity of studied bacteriocin is bactericidal. Obtained results suggest that bacteriocin producing strain *P. pentosaceus* ST0408, evaluated in current study can be considered safe and expressed bacteriocin can be implemented




IAFP Latino 2024

Simpósio Latinoamericano
em Segurança dos Alimentos

Santos - SP - Brasil
11 a 14 Nov, 2024

for the control of *L. monocytogenes* in the fermented dairy products.

Agradecimentos: Acknowledgement: To FCT (UIDB/05937/2020 and UIDP/05937/2020), Fundação para a Ciência e a Tecnologia, Portugal; FAPESP, processos 2023/05394-9; 2023/14944-2, Sao Paulo, SP, Brazil.



BRAFP



International Association for
Food Protection