

Listeria monocytogenes biofilm prevention using *Enterococcus faecium* semi-purified bacteriocin

João Marcos Scafuro Lima^{1,2}, Uelinton Manoel Pinto¹, **SVETOSLAV DIMITROV TODOROV**^{1,2,3}

¹. Faculty of Pharmaceutical Sciences, University of São Paulo, Food Research Center, Sao Paulo, Brazil

². Faculty of Pharmaceutical Sciences, University of São Paulo, ProBacLab, Laboratório de Microbiologia de Alimentos, Departamento de Alimentos e Nutrição Experimental, Sao Paulo, Brazil

³. Instituto Politécnico de Viana do Castelo, CISAS - Center for Research and Development in Agrifood Systems and Sustainability, Viana Do Castelo, Portugal

Listeria monocytogenes is an opportunistic foodborne pathogen, a ubiquitous bacterium able to persist and adhere in food processing environment forming biofilms. However, lactic acid bacteria, beyond their beneficial applications, can produce an arsenal of antimicrobials, including bacteriocins. The aim of this study is to evaluate the prevention of biofilm formation of different serovars of *L. monocytogenes* by applying a semi-purified bacteriocin produced by *Enterococcus faecium* ST02JL, a strain isolated from artisanal cheesemaking environment. *E. faecium* ST02JL was cultured in MRS broth for 18h at 37°C, and cell-free supernatant (CFS) was obtained by centrifugation (4000×g, 30 min, 4°C), filter sterilized (0.22 µm), and heat-treated (80°C for 10 min). Antimicrobial proteins were precipitated (60% ammonium sulfate saturation), collected by centrifugation (20000×g, 60 min, 4°C), and re-suspended in 25 mM phosphate buffer (pH 6.5). Chromatography on SepPakC18 column was applied for further separation and targeted proteins eluted by 60% iso-propanol/25 mM phosphate buffer. For the biofilm inhibition study, 18h-old cultures of the studied *L. monocytogenes* strains were grown in TSBYE to a final cell concentration of ~107 CFU/mL. Each strain was transferred to 96-well microtiter plates and supplemented with semi-purified bacteriocin at different levels of activity. The biofilm challenge assay was performed at 25°C for 48h in triplicates, where quantification of biofilms was made through Crystal Violet Assay. The semi-purified bacteriocin produced by *E. faecium* ST02JL strain promotes the biofilm inhibition of each evaluated *L. monocytogenes* strain in the treatment of at least 800 AU/mL bacteriocin. Considering that, a previous estimated MIC of ≥3200 AU/mL for the bacteriocin produced by *E. faecium* ST02JL, our observations confirmed the even sub-MIC bacteriocin activity has inhibitory effects against *Listeria monocytogenes* biofilm formation. Bacteriocins can be an alternative for controlling the biofilm formation in the industry surfaces.

Agradecimentos: Acknowledgement: To FAPESP (2013/07914-8, 2023/05394-9), Brazil.

