


Production of Pectinases from *Bacillus* sp. SMIA-2 for Hydrolyzing Pectin from Passion Fruit Flour to Improve the Colonic In Vitro Fermentation

Erica Cruz¹, Larissa Pacheco Ferreira², Débora Preceliano de Oliveira¹, Meire Lelis Leal Martins², João Paulo Fabi¹

¹. University of São Paulo, Department of Food Science and Experimental Nutrition / Faculty of Pharmaceutical Sciences, São Paulo/SP, Brazil

². State University of North Fluminense, Food Technology Laboratory, Campos dos Goytacazes/RJ, Brazil

Pectin is a structural heteropolysaccharide found in the cell walls of terrestrial plants. Pectin can be hydrolyzed into smaller molecules through chemical, thermal, enzymatic, or microbial processes. The hydrolysis breaks down the complex polysaccharides into simpler pectic oligosaccharides. Enzymatic hydrolysis, using pectinases, is a method for producing pectic oligosaccharides with specific properties and functionalities. This work aimed to produce satisfactory levels of pectinases by *Bacillus* sp. SMIA-2 in submerged cultures containing agro-industrial residues as substrates to hydrolyze pectin from passion fruit peel flour, evaluating the resulting product (partially hydrolyzed pectin) regarding its physicochemical properties and improving the colonic in vitro fermentation. *Bacillus* sp. SMIA-2 secreted pectinases (18.43 U/mL), avicelase (0.0487 U/mL), carboxymethylcellulase (0.0405 U/mL), amylase (0.1365 U/mL), xylanase (0.0220 U/mL) and protease (6.3000 U/mL) in submerged cultures containing passion fruit peel flour and corn steeping water. Pectinases were the enzymes that showed the greatest enzymatic activity. In this way, it was possible to create, unprecedentedly, a series of methodologies for producing hydrolytic enzymes - especially proteolytic and glycoside enzymes - that are highly sustainable and low-cost by using waste from the food industry and very resistant and propagating bacteria. Different enzymes are required for the degradation of complex substrates present in the plant cell wall. The bacterial strain used in this study was a thermophilic *Bacillus* sp. SMIA-2 isolated from a soil sample collected in Campos dos Goytacazes city, Rio de Janeiro, Brazil. The method for in vitro fermentation was based on studies by Jonathan et al. (2012) and Williams et al. (2005), in which the inoculum was immediately prepared after receiving the fecal material. The feces were pooled and diluted in a sterile anaerobic solution of 0.9% NaCl and 20% glycerol, in a 1:6 (w/v) ratio. The material was homogenized and filtered (Miracloth, EMD Millipore), and the inoculum was placed in a CO₂ saturation container to maintain the anaerobic condition. The prepared fermentation medium was the basal solution (76%), vitamin/phosphate buffer solution (1%), bicarbonate buffer (1%), and reducing agent (4%). All materials were sterilized, and the medium and inoculum flasks were continuously bubbled with CO₂. The carbohydrates used as substrate were modified passion fruit flour, unmodified passion fruit flour, FOS, and polydextrose. In each fermentation bottle, 200 mg of a specific carbohydrate, 24 mL of fermentation medium, and 1.5 mL of the inoculum were added. The environment was saturated with CO₂ in bottles with screw caps, and the samples were incubated in a water bath at 37°C. The test for each treatment was done in triplicate for analysis at each fermentation time (0 h, 8 h, 16 h, and 24 h). At each point in the fermentation time, pH and pressure were evaluated to indicate that fermentation had occurred. During fermentation, the production of organic acids caused a




IAFP Latino 2024

Simpósio Latinoamericano
em Segurança dos Alimentos
Santos - SP - Brasil
11 a 14 Nov, 2024

decrease in the pH of the culture medium. Regular pH monitoring indicated that fermentation had occurred. Other analyses such as the fecal microbiota composition via 16S rDNA sequencing will be carried out to understand the ecosystem and the role of our substrate.

Agradecimentos: The authors are thankful for the financial support granted by the CAPES, Fapesp, PRIP-USP, UENF.



BRAFP



International Association for
Food Protection