

Genomic analysis of Avian pathogenic *Escherichia coli* (APEC) isolated from a Brazilian slaughterhouse

**Nayla Kellen de Oliveira Ventura**<sup>1</sup>, Jordano Alexandre de Carvalho<sup>1</sup>, Rafaela de Melo Tavares<sup>1</sup>, Rodrigo Sebastião Machado de Freitas<sup>1</sup>, Fernanda Simone Marks<sup>1</sup>, Luís Augusto Nero<sup>1</sup>, Ricardo Seiti Yamatogi<sup>1</sup>

<sup>1</sup> Universidade Federal de Viçosa, Viçosa / Minas Gerais, Brasil

Avian pathogenic *Escherichia coli* (APEC) is a pathogen that can cause extraintestinal infections responsible for many economic losses in the poultry food chain. Besides, APEC is also presumed to be a zoonotic pathogen, causing health risks to humans. This work aimed to provide a genomic portrait of APEC isolated from broiler chickens from a Brazilian slaughterhouse in Minas Gerais state. The Whole Genome Sequencing analysis was conducted using software for trimming, assembly, annotation, and virulence factors/antimicrobial resistance investigation (in order Trimmomatic v0.39, SPAdes v3.15.5, Prokka v1.13.3, BV-BRC v.3.35.5, CARD v3.2.9). Also, the isolates were screening for phylogroup (ClermoTyping), serotype (ECTyper - 1.0.0), and sequence type (BV-BRC v.3.35.5). A total of 35 isolates were characterized as APEC by the presence of at least 2 of the 5 virulence genes mostly associated with this pathotype (*iroN*, *ompT*, *hlyF*, *iutA* and *iss*). The most prevalent phylogroups were B1 (15/35), D and F (4/35 each). Most of the APEC belong to serogroup O9 (6/35). Sequence types ST155 (7/35), ST69, ST361, ST3258, and ST5459 (2/35 each) were the most identified. The highest levels of antimicrobial resistance were detected in the peptide and fluoroquinolone; macrolide, and penam classes. Peptides showed *bacA*, *eptA*, *PmrF*, *OmpA* and *YojI* as frequent genes. Meanwhile fluoroquinolone; macrolide; and penam revealed *gadX*, *gadW*, *CRP*, *mdtE* and *mdtF* as predominant. Additionally, beta-lactamase genes, including *bla*TEM, *bla*CTX-M, *bla*SHV, and *ampC* were also identified, underscoring the concern given to the beta-lactam antibiotics as a human treatment for various diseases. This study provides a better characterization of this pathotype found in the Minas Gerais state, and contributes significantly to understanding the characteristics, circulation, and resistance genes among these isolates.

**Agradecimentos:** CAPES, CNPq, FAPEMIG.

