Simpósio Latinoamericano em Segurança dos Alimentos Santos - SP - Brasil 11 a 14 Nov, 2024

Evaluation of Salmonella enterica behavior in hummus under different temperature conditions

**Alice Karine da Silva**<sup>1</sup>, Jean Carlos Correia Peres Costa<sup>2</sup>, Jaqueline Barbosa da Silva<sup>1</sup>, Inayara Cristina Alves Lacerda<sup>1</sup>, Verônica Ortiz Alvarenga<sup>3</sup>

<sup>1.</sup> Universidade Federal de Minas Gerais, Campus Pampulha- Faculdade de Farmácia, Belo Horizonte/Minas Gerais, Brasil

<sup>2.</sup> University of Lleida, Department of Food Technology, Engineering and Science, Lleida, Spain

<sup>3.</sup> Universidade Federal de Minas Gerais, Campus Pampulha- Faculdade de Farmácia, Belo Horizonte/Minas Gerais, Brasil

Hummus is a ready-to-eat (RTE) food made from cooked chickpeas, olive oil, lemon juice, tahini, salt, and water, and has been widely consumed worldwide. However, during the production process, contamination by pathogenic microorganisms such as Salmonella enterica can occur. Therefore, this study assessed the behavior of three serotypes of Salmonella enterica in hummus paste stored at different temperatures. The hummus was prepared under aseptic conditions, following a traditional formulation. After preparation, three strains of Salmonella (S. Senftemberg, S. Typhimurium, and S. Enteritidis) were inoculated individually into the hummus at ca. of 3 log CFU/g and then stored at 4, 7, 10, 12 and 15 °C for 10 days. Potential growth ( $\delta$ ) was calculated for each of the *S. enterica* strains through the difference between the maximum observed and initial counts. Additionally, the Baranyi and Roberts (BAR) model was used to obtain the kinetic parameters of growth, including lag time ( $\lambda$ ; h), maximum specific growth rate ( $\mu_{max}$ ;  $h^{-1}$ ), and maximum population density ( $N_{max}$ ; log CFU/g), while the Weibull model was used to obtain the time for first decimal reduction (h). Both models were applied to each of the three strains of Salmonella. The  $\delta$  values ranged from 6.17 to -1.33, 6.28 to -2.88, and 6.07 to -1.74 log CFU/g for S. Enteritidis, S. Typhimurium, and S. Senftemberg, respectively, within the studied temperature range. The estimated  $\mu_{max}$ for the Salmonella enterica strains in hummus at 10-15 °C, fitted by the BAR model, ranged from 0.013 to 0.205 h<sup>-1</sup>. The  $\lambda$  for the pathogen strains stored within the temperature range varied from 9 to 98 h. However, at 12  $^{\circ}$ C, no  $\lambda$  was observed for any strains. The S. Senftemberg was the fastest strain, with  $\mu_{max}$ = 0.205 h<sup>-1</sup> when the product was stored at 15  $^{\circ}$ C, meanwhile the lowest  $\mu_{max}$  was observed for the same serotype when stored at 10 °C, with a  $\mu_{max} = 0.013 \text{ h}^{-1}$ . All tested strains achieved N<sub>max</sub> values >7.0 log CFU/g at temperatures ranging from 10 to 15 °C. At 10 °C, the S. Enteritidis strain exhibited the highest  $\mu_{max}$  of 0.074 h<sup>-1</sup> and longest  $\lambda$ , which was 98 h. For storage temperatures of 4 and 7 °C, the pathogen\'s survival in hummus can be observed, resulting in decimal reduction times ranging from 15.9 to 167.5 h. The highest value for decimal reduction corresponded to the S. Senftemberg strain (167.5 h) when the hummus was stored at 7 °C. However, the smallest reduction was observed in the S. Typhimurium (15.9 h) strain when stored at 4 °C. At refrigeration, abuse temperatures (15 and 12 °C) and at 10 °C, the growth of S. enterica in hummus increased. Meanwhile, at lower storage temperatures (4 and 7 °C), S. enterica survived in the food. The estimated parameters can be useful in assessing the shelf life of product and establishing measures to control the growth of the pathogen.

BRAFF

International Association for

Food Protection



**Agradecimentos:** Órgãos Financiadores: Conselho Nacional de Desenvolvimento Científico e Tecnológico-CNPQ (Processo 160795/2022-2; Processo 311863/2022-1), Capes Código 001.

## BRAFP